

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
11 October 2001 (11.10.2001)

(10) International Publication Number  
**PCT WO 01/75088 A1**

- (51) International Patent Classification<sup>7</sup>: C12N 9/64, C07K 14/47, C12N 9/26 (74) Agent: E. BLUM & CO.; Vorderberg 11, CH-8044 Zürich (CH).
- (21) International Application Number: PCT/IB00/00552 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 1 May 2000 (01.05.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/194,609 5 April 2000 (05.04.2000) US (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): **ESBAT-ECH AG** [CH/CH]; Winterthurerstrasse 190, CH-8057 Zürich (CH).


- (72) Inventors; and  
(75) Inventors/Applicants (*for US only*): **LUETHI, Urs** [CH/CH]; Handgasse 2, CH-8180 Bülach (CH). **BARBERIS, Alcide** [CH/CH]; Allenmoosstrasse 64, CH-8057 Zürich (CH).

Published:  
— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD FOR IDENTIFY POLYPEPTIDES WITH PROTEASE ACTIVITY

### SUC2-APP fusion with N-terminal truncated APP



(57) Abstract: The present invention describes methods for the identification of proteins with protease activity and methods for the analysis of protein-protein interactions of membrane proteins.

BEST AVAILABLE COPY

WO 01/75088 A1

## METHOD FOR IDENTIFY POLYPEPTIDES WITH PROTEASE ACTIVITY

Technical Field

The present invention relates to methods for  
5 the identification of proteins with secretase activity  
and methods for the analysis of interactions between  
membrane proteins.

Background Art

10 Protein secretion is central to the proper  
development and function of eucaryotic organisms.  
Moreover, several pathophysiological processes such as  
neurodegeneration, oncogenesis, apoptosis and  
inflammation are associated with the malfunction or  
15 aberrant regulation of protein secretion. It has become  
clear that there is no single biosynthetic mechanism  
common to all secretory proteins. Secretion of proteins  
can occur through either the regulated or constitutive  
pathways and, in some cell types, this secretion can be  
20 polarized to distinct cellular domains. An increasing  
number of proteins are now recognized as being derived  
from integral membrane proteins of type I and type II  
topology and, in this case, the secretory event involves  
their selective post-translational hydrolysis from the  
25 cell surface. This secretion is catalyzed by proteases  
known as secretases. The cleavage of membrane proteins  
generally occurs near the extracellular face of the  
membrane, although in some cases it has been shown also  
to occur within the transmembrane domain. Proteins  
30 secreted in this fashion include membrane receptors and  
receptor ligands, ectoenzymes, cell adhesion molecules  
and others. Examples of protein secretion through the  
action of secretases include the vasoregulatory enzyme  
ACE (angiotensin converting enzyme), the tumor necrosis  
35 factor (TNF) ligand and receptor superfamilies, the  
transforming growth factor- $\alpha$ , certain cytokine receptors,  
the Alzheimer's amyloid precursor protein (APP) and

others (Hooper, N.M., Karran, E.H., and Turner, A.J. (1997) Biochem. J. 321, 265-279). Most secretases have so far eluded identification and cloning. Evidently, the cloning of the genes encoding these enzymes would  
5 dramatically facilitate their characterization and provide the unique condition to design specific inhibitors or stimulators that target discrete secretases.

In the past, different methods have been used  
10 to clone genes encoding secretases. The most widely used method has been purification of the protein and subsequent isolation of the corresponding cDNA sequence (Black, R.A et al., (1997) Nature 385, 729-733; Moss M.L. et al., (1997) Nature 385, 733-736; Howard L. et al.,  
15 (1996) Biochem. J. 317, 45-50)). An alternative method has been used by Yan et al., who have cloned a APP  $\beta$ -secretase by exploiting pharmacological data, which allowed classification of this secretase as an aspartyl protease, to scan the almost completely sequenced genome  
20 of the nematode worm *C. elegans* for candidate protease genes, and to subsequently use such candidates to identify human homologues ( Yan R. et al., (1999) Nature 402, 533-537). A third method has been applied by Vassar et al., who have cloned the same  $\beta$ -secretase gene by  
25 screening pools of human cDNAs which were expressed in a mammalian cell line and analyzed for increased release of the APP  $\beta$  proteolytic product (Vassar R. et al., (1999) Science 286, 735-741).

It is already known from Hawkins et. al.,  
30 (Hawkins C.J., Wang S.L., and Hay B.A., (1999) Proc.Natl.Acad.Sci.USA 96, 2885-2890) to use yeast for screening for cytoplasmic proteases of the group of caspases and their regulators. In the described method a fusion protein is created in which a transcription factor  
35 is linked to the intracellular domain of a transmembrane protein by caspase cleavage sites. The transcription factor is part of a reporter system thus that in the

presence of a caspase the transcription factor is released and induces transcription of a reporter gene and thus allows identification of positive cells. The method allows only the isolation of members of the caspase group.

In the past various methods were developed to identify protein-protein interactions in vivo. The two hybrid system is a very powerful tool for the in vivo analysis of interactions between soluble proteins (Bartel P.L., and Fields S., (1995) Methods Enzymol., 254, 241-263). The split-ubiquitin system is an alternative method for the analysis of interactions between soluble proteins (Johnsson N., and Varshavsky A., (1994) Proc.Natl.Acad.Sci.USA, 91, 10340-10344). Stagljar et al., (Stagljar I, Korostensky C., Johnsson N., and Te Heesen S., (1998) Proc.Natl.Acad.Sci.USA, 95, 5187-5192) describe a genetic system based on split-ubiquitin for the analysis of interactions between membrane proteins in vivo. This method provides a potentially useful tool for screening of interaction between membrane proteins.

The importance of secretases in many biological processes makes them attractive novel targets for the development of therapeutic drugs and this raises the need for reliable screening systems for the cloning of new members of the secretase family which are more sensitive and efficient than the known traditional methods.

#### Disclosure of the Invention

Hence, it is a general object of the present invention to provide a method for the identification of a secretase wherein suitable host cells are transformed to express under suitable conditions a target membrane protein as a fusion protein with a secreted protein and said suitable host cells transformed to express said fusion protein are further transformed with a library encoding candidate secretases and said host cells

transformed with said fusion protein and said library are cultivated under conditions allowing expression of both said fusion protein and said library and said conditions allowing cell survival only in presence of said secreted  
5 protein that has been released from said fusion protein by a secretase encoded by said library.

The present invention furthermore provides a secretase that can be obtained by a method of the invention. The term secretase as used herein should be  
10 understood to include all types of proteins with protease activity.

Another object of the present invention relates to a method for the identification of a membrane protein that is a substrate of a known secretase wherein  
15 suitable host cells are transformed to express under suitable conditions said secretase and said host cells transformed to express said secretase are further transformed with a library encoding fusion proteins with a secreted protein and a candidate substrate and said  
20 host cells are cultivated under conditions allowing expression of both the secretase and the library encoded fusion protein and allowing cell survival only in the presence of said secreted protein that has been released from said fusion protein by said secretase due to the  
25 interaction of said secretase with said library encoded substrate.

Another object of the present invention concerns a method for screening for a protein interacting with a target membrane protein wherein suitable host  
30 cells are transformed to express under suitable conditions said membrane protein as a fusion protein with a secreted protein and the two moieties of said fusion protein are linked by a recognition sequence that is cleaved by a defined secretase. Said host cells  
35 transformed to express said fusion protein are further transformed with a library encoding fusion proteins with said defined secretase and candidate interaction partners

of said known membrane protein and cultivated under conditions allowing expression of both the fusion protein and the library and allowing cell survival only in the presence of an interaction of the known membrane protein with the library encoded interaction partner protein. Due to the interaction of said known membrane protein and the library encoded partner protein the secretase of said library fusion protein is brought into vicinity of said secretase recognition sequence linking the target membrane protein and the secreted protein and this vicinity allows the release of the secreted protein from said fusion protein and thus cell survival.

Appropriate culturing conditions of the cells must be used such that said fusion protein is not efficiently cleaved by the secretase in the absence of a protein-protein interaction between said known membrane protein and said library encoded partner protein.

In a preferred embodiment said secretase is modified, preferably said modification is such that the membrane-anchoring domain of said secretase is deleted.

This method allows the identification of interactions between membrane bound proteins or between ER/Golgi luminal proteins.

The invention furthermore provides an interacting protein that can be obtained by a method of the invention.

Another object of the present invention is a method for screening for a secretase wherein suitable host cells are transformed to express under suitable conditions a target membrane protein as a fusion protein comprising the known membrane protein, a secreted protein and a transcriptional activator which is part of a reporter system that is stably integrated into the genome of said host cells. Said host cells are further transformed with a library encoding candidate secretases and cultured under conditions allowing expression of both said fusion protein and said library and allowing cell

survival and selection of positive cells only in the presence of a library encoded secretase that cleaves said fusion protein and thus releases said secreted protein and the transcription activator from said fusion protein  
5 thus allowing cell survival and activation of a reporter gene and thus selection of positive cells.

The presence of a secreted protein and a transcriptional activator in a fusion protein with a target transmembrane protein allows the rapid  
10 identification of secretases cleaving a transmembrane protein in its transmembrane region. In such a case both the secreted protein is released from the fusion protein and can be secreted thus allowing cell survival and the transcriptional activator is released and can enter the  
15 cell nucleus and induce transcription from the reporter gene construct and thus allowing selection of positive cells.

Protein-protein interactions can, preferably, be monitored by a transcriptional readout, preferably by  
20 the expression of the HIS3 gene whose expression is under control of said transcriptional activator. Many transcriptional activators useful for the described methods are well known to the skilled person and are for example LexA-VP16.

25 The invention provides furthermore a secretase that can be obtained by a method of the invention.

Secretases are generally expressed as pro-enzymes which are dependent on processing to be  
30 activated. The need of an external processing activity exists when the prosecretase is not capable of autoproducting. Therefore the present invention provides a method for the identification of a protein that is able to process a specific secretase wherein suitable host  
35 cells are manipulated to express under suitable conditions a target membrane protein as a fusion protein with a secreted molecule and a unprocessed specific

secretase. Said cells are further transformed with a library encoding candidate processing proteins of said secretase and cultivated under conditions allowing expression of said fusion protein, said unprocessed  
5 specific secretase and said library and said conditions furthermore allowing cell survival only in the presence of said secreted protein that has been released from said fusion protein by said secretase that has been processed by the library encoded processing protein.

10 In a preferred embodiment of all the above mentioned objects of this invention the host cell is an eucaryotic cell, in particular a yeast cell. The target membrane protein preferably is a transmembrane protein, preferably a type I or a type II transmembrane protein.  
15 The term membrane protein as used herein comprises full length proteins as well as fragments thereof. Depending on the particular topology of the membrane protein the secreted protein moiety is fused to the N-terminus or the C-terminus of said protein, such that said moiety faces  
20 the ER lumen. General molecular biological and biochemical methods well known to the person skilled in the art are applied to determine the most suitable fusion protein of the target membrane protein and the secreted protein to result in a functional secreted protein after  
25 secretase cleavage (Maniatis et al., Molecular Cloning: A Laboratory Manual, New York: Cold Spring Harbor Laboratory, 1989). There are many strategies well known in the technical field to construct DNA libraries appropriate for the purposes of the present invention and  
30 to clone them in suitable vectors for expression in host cells (Maniatis et al., Molecular Cloning: A Laboratory Manual, New York: Cold Spring Harbor Laboratory, 1989).

The introduction of the expression constructs of this invention into said suitable host cells can be  
35 performed by cotransformation or more preferably by sequential transformation, wherein in a much preferred embodiment of this invention said host cells are first



transformed with the expression construct encoding said target protein or a fusion protein of said target protein and in a second transformation step said cells transformed to express said target protein or a fusion  
5 protein of said target protein are transformed with the second expression construct. The second construct can be a library encoding fusion proteins or a library encoding single proteins.

In a preferred embodiment of the present  
10 invention the secreted protein is an protein with invertase activity or functional fragments of a protein with invertase activity, preferably a yeast invertase or functional fragments of a yeast invertase. Yet it is obvious for the man skilled in the art that other  
15 secreted proteins e.g the yeast pheromone  $\alpha$ -peptide, can be used in the scope of the present invention as a reporter system.

Other suitable host cells for all the objects of this invention are tissue culture cell lines whose  
20 growth is dependent on a protein growth factor. These cells can be manipulated such that they express in one embodiment of this invention the growth factor fused to a target membrane protein. Since the modified growth factor can not be secreted, these cells are dependent on a  
25 exogenous growth factor. Said host cells are then further transfected with a library and cultivated under conditions allowing the expression of both said fusion protein of the target membrane protein with said growth factor and the library and said conditions furthermore  
30 allowing growth of said host cells only in the presence of said growth factor that has been released from said fusion protein by a library encoded secretase. A suitable host cell line is for example PC12 whose growth is depending on the presence of NGF (nerve growth factor).

35

#### Brief Description of the Drawings

The invention will be better understood and objects other than those set forth above will become apparent when consideration is given to the following detailed description thereof. Such description makes reference to the annexed drawings, showing the following:

Fig 1 A: Deletion strategy of the YAP3 gene of *S.cerevisiae*.

Fig 1 B: Confirmation of homologous recombination at the YAP3 locus by PCR genotyping.

Fig 1 C: Deletion strategy of the MKC7 gene of *S.cerevisiae*.

Fig 1 D: Confirmation of homologous recombination at the MKC7 locus by PCR genotyping.

Fig 2 A: Deletion strategy of the SUC2 gene in *S. cerevisiae*.

Fig 2 B: Confirmation of homologous recombination at the SUC2 locus by PCR genotyping.

Fig 3: Genetic test for *suc2* disruption

Fig 4 A: Western blot analysis of SUC2 fusion with N-terminal truncated APP.

Fig 4 B: Colony formation of ULY 2 cells expressing SUC2-APP fusion proteins on sucrose plates.

Fig 5: SUC2-APP(590-695) is activated in  $\alpha^+$  background.

Fig 6: ER retrieval signal in  $\alpha^+$  and  $\alpha^-$  background.

#### Modes for Carrying Out the Invention

The ability of yeast such as *Saccharomyces cerevisiae* to utilize sucrose as a carbon source depends on the secretion of the enzyme invertase, which cleaves sucrose to yield glucose and fructose ( Carlson, M. et al. (1983) Mol. Cell. Biol. 3, 439-447). Indeed, deletion of the entire SUC2 gene, which encodes the invertase protein, or deletion of the signal peptide, which prevents secretion of the invertase, cripples the ability of yeast cells to grow on sucrose medium ( Perlman, D.,

and Halvorson, H.O. (1981) *Cell* 25, 525-536; Carlson, M., and Botstein, D. (1982) *Cell* 28, 145-154; Kaiser, C.A., and Botstein, D. (1986) *Mol. Cell. Biol.* 6, 2382-2391). Thus, the manipulation of invertase expression and  
5 secretion provides a convenient genetic selection for engineered yeast cells, as it has been shown by two research groups who have utilized this growth selection system to isolate human cDNAs encoding secreted proteins ( Klein, R.D. et al. (1996) *Proc. Natl. Acad. Sci. USA*  
10 93, 7108-7113; Jacobs, K.A. et al. (1997) *Gene* 198, 289-296; patents: US005536637A, US00571211A, WO 97/40146). In these works, the authors have used yeast strains deleted for the endogenous SUC2 gene to express human cDNAs fused to a modified SUC2 gene lacking its leader sequence,  
15 which encodes the secretion signal sequence. Heterologous secreted proteins appropriately fused to the N-terminus of the modified, non-secreted invertase could be identified through positive selection because they provided the necessary signals to restore invertase  
20 secretion, thus restoring cell growth on sucrose medium.

Expression and localization of the invertase protein have also been monitored by alternative techniques such as colorimetric ( Goldstein, A., and Lampen, O.J. (1975) *Methods Enzymol.* 42, 504-511) and  
25 immunodetection assays. For example, these techniques have been used to identify protein-sorting sequences that mediate localization to yeast mitochondria ( Emr, S.D. et al. (1986) *J. Cell. Biol.* 102, 523-533), vacuoles ( Klionsky, D.J. et al. (1988) *Mol. Cell. Biol.* 8, 2105-  
30 2116; Tague, B.W. et al. (1990) *Plant Cell* 2, 533-546 ), or endoplasmic reticulum ( Gaynor, E.C. et al. (1994) *J. Cell. Biol.* 127, 653-665; Boehm, J. et al. (1994) *EMBO J.* 13, 3696-3710), to determine the effects of mutations within signal sequences ( Ngsee, J.K. et al. (1989) *Mol.*  
35 *Cell. Biol.* 9, 3400-3410), and to monitor the amounts of human proteins expressed from yeast ( Hitzeman, R.A. et al. (1990) *Methods Enzymol.* 185, 421-440).

In one embodiment of the present invention, the identification of a secretase activity expressed in yeast is based on its ability to cleave a specific target membrane protein fused to the invertase enzyme. Because  
5 of its fusion with the membrane-bound protein, this invertase is not secreted; consequently, these yeast cells, which lack the endogenous invertase, cannot grow on sucrose medium. However, in the presence of a secretase activity that specifically recognizes and  
10 cleaves the membrane-bound protein, the invertase enzyme is liberated from its anchor and it is secreted to the periplasm where it can hydrolyze sucrose, thus allowing cell growth on sucrose medium.

For the experiments described here,  
15 chimerical proteins were used bearing the invertase enzyme fused to different portions of the membrane-bound Amyloid- $\beta$  Precursor Protein (APP). In human cells, APP, a type I transmembrane protein, can be processed by three types of proteases denoted  $\alpha$ ,  $\beta$ - and  $\gamma$ -secretases.  
20 Cleavage by the latter two generates the 40 and 42 amino acids A $\beta$  peptides involved in Alzheimer's disease, while the  $\alpha$ -secretase cleaves APP near the middle of the A $\beta$  sequence. Cleavage of APP by  $\alpha$  and  $\beta$ -secretases occurs at the luminal/extracellular face of the membrane, whereas  
25 cleavage by  $\gamma$ -secretase has been shown to occur within the transmembrane domain of APP. APP expressed in the yeast *Saccharomyces cerevisiae* is efficiently processed by endogenous proteases which cleave the protein almost exclusively at the  $\alpha$ -site. Two research groups have  
30 independently shown that the glycosyl-phosphatidylinositol-linked aspartyl proteases Yap3p and Mkc7p are primarily responsible for  $\alpha$ -secretase-type cleavage of APP, which results in release and secretion of soluble APP into the periplasm ( Zhang, W. et al.  
35 (1997) Biochim. Biophys. Acta 1359, 110-122; Komano, H. et al. (1998) J. Biol. Chem. 273,31648-31651). Deletion of YAP3 and MKC7 in a vacuolar protease-deficient strain

abolished  $\alpha$ -secretase cleavage, which could be restored by reintroducing MKC7 or YAP3 on single copy plasmids (Komano, H. et al. (1998) J. Biol. Chem. 273,31648-31651). Here we present results showing that a yeast  
5 strain deleted for the endogenous SUC2 gene and expressing the invertase enzyme fused to the membrane-bound APP can efficiently grow on sucrose only when the Yap3p and Mkc7p secretases that cleave APP at the  $\alpha$ -site are co-expressed with the fusion protein.

10 The experiments utilize two *Saccharomyces cerevisiae* strains, both of which have been deleted for the SUC2 gene, and one of them has been additionally deleted for the YAP3 and MKC7 genes. Fig. 1 shows the strategy that was used to delete the YAP3 and MKC7 genes.  
15 For the YAP3 knock-out, a kanamycin resistance cassette specifically constructed for yeast expression (Steiner, S., and Philippsen, P. (1994) Mol. Gen. Genet. 242, 263-271) was amplified by PCR using primers that possess at their 5' termini sequences homologous to YAP3 regions  
20 (Fig. 1A). The yeast strain JPY9 (Barberis, A. et al. (1995) Cell 81, 359-368) was transformed with this PCR product and plated on a selective agar medium containing the kanamycin analog G418. Only those cells that have steadily integrated the kanamycin resistance cassette  
25 could grow and form colonies on these selective plates. In order to check whether integration had occurred in a homologous manner, namely at the YAP3 locus, we performed genotyping by PCR using two sets of primers (Fig. 1B). The yap3 derivative of JPY9 was named STY1. To knock out  
30 the MKC7 gene in STY1, a functional HIS3 gene-cassette was amplified by PCR using primers that bear at their 5' termini sequences homologous to the MKC7 gene (Fig. 1C). STY1, which lacks the endogenous HIS3 gene, was transformed with this PCR product and plated on a  
35 selective agar medium lacking histidine. Only those cells that have steadily integrated the HIS3 gene-cassette could grow and form colonies on these selective plates.

In order to check whether integration had occurred in a homologous manner, namely at the MKC7 locus, we performed genotyping by PCR using three sets of primers (Fig. 1D). The yap3, mkc7 double knock-out strain was named STY2.

5 Fig. 2 shows the strategy we used to knock out the SUC2 gene in both STY1 and STY2 strains. A functional URA3 gene-cassette was amplified by PCR using primers that bear at their 5' termini sequences homologous to the SUC2 gene (Fig. 2A). The yeast strains JPY9 and STY2, which  
10 lack a functional URA3 gene, were transformed with this PCR product and plated on a selective agar medium lacking uracil. Only those cells that have steadily integrated the URA3 gene-cassette could grow and form colonies on these selective plates. In order to check whether  
15 integration had occurred in a homologous manner, namely at the SUC2 locus, we performed genotyping by PCR using two sets of primers (Fig. 2B). The suc2 derivative of JPY9 was named ULY1, while the suc2 derivative of STY2 was named ULY2.

20 The phenotype of the suc2 strains was tested as follows. The SUC2 gene was cloned on a single copy yeast expression vector and introduced into the strains ULY1 and ULY2. Growth of the transformed cells on sucrose plates was monitored at different time points and  
25 compared to growth of the same strains that were transformed with a vector that did not bear the SUC2 gene. Fig. 3 shows that, while growth on sucrose plates of the strains lacking SUC2 was dramatically impaired, expression of the vector-borne SUC2 gene complemented the  
30 suc2 phenotype. The growth rate of the complemented strains was indistinguishable from that of wild type cells (data not shown). Moreover, no significant growth difference was observed between ULY1 and ULY2 in this experiment.

35 The invertase-APP fusion protein tested in the experiments described here is composed of full-length invertase (amino acid residues 1-532) sequentially

followed by a 9 amino acid HA-tag and by a C-terminal portion of APP (amino acid residues 590-695), which contains, in addition to the transmembrane domain (aa 622-646), the cleavage sites for all three mammalian  
5 secretase ( $\beta$  between aa 596 and 597,  $\alpha$  between aa 612 and 613, and  $\gamma$  between aa 636 and 637 or between aa 638 and 639). It has been shown that an APP deletion mutant containing amino acid residues 590 to 695 is proteolytically processed in a proper manner in mammalian  
10 cells (Citron, M. et al. (1995) Neuron 14, 661-670). In order to control and predict the activity of the invertase-HA-APP fusion which is expected to be secreted upon cleavage by the  $\alpha$ -secretases in yeast, we constructed an invertase-HA-APP fusion protein bearing an  
15 APP sequence ending at the  $\alpha$  site (aa 590-612). Expression of both fusion proteins in the strain ULY2, which lacks  $\alpha$ -secretase activities, was confirmed by Western blot analysis using anti-HA antibodies (Fig. 4A). Fig. 4B shows the effect that these fusion proteins had  
20 on the rate of ULY2 cell growth and colony formation on sucrose plates. As a positive control, the SUC2 expression plasmid described above was used; an empty vector, i.e. not expressing any SUC2 gene, was used as a negative control. Growth on sucrose plates of ULY2 cells  
25 expressing either the wild type SUC2 gene or the  $\alpha$ -truncated invertase-HA-APP (590-612) fusion protein was indistinguishable (Fig. 4B, top two rows), indicating that fusion of this portion of APP to the C-terminus of the invertase did not influence invertase function. On  
30 the other hand, expression of the invertase-HA-APP (590-695) fusion protein, which bears the transmembrane domain, visibly reduced growth of the transformed ULY2 cells, although not to the extent observed with cells transformed with the empty vector (Fig. 4B, bottom two  
35 rows).

In order to test the hypothesis that co-expression of the yeast  $\alpha$ -secretases Yap3p and Mkc7p with

the invertase-HA-APP(590-695) fusion protein would cause cleavage of the APP sequence at the  $\alpha$  site, thus allowing efficient secretion of the invertase, we compared growth on sucrose plates of ULY1 (containing the  $\alpha$ -secretases;  $\alpha$  +) with ULY2 (lacking the  $\alpha$ -secretases;  $\alpha$ -) after transformation with the vector expressing the invertase-HA-APP(590-695) fusion protein. Fig. 5 shows that co-expression of active  $\alpha$ -secretases with this invertase fusion protein in yeast cells lacking endogenous invertase restored their ability to grow on sucrose plates to an extent very similar to the positive control, in which cells were transformed with a vector expressing wild type invertase. In contrast, yeast cells that do not co-express the  $\alpha$ -secretases with the invertase fusion protein have a reduced growth rate on sucrose plates, although, as previously shown, not to the extent observed with cells transformed with the empty vector.

One explanation for the observed residual activity of the invertase-HA-APP (590-695) fusion protein expressed in cells lacking Yap3 and Mkc7  $\alpha$ -secretases is that this membrane-bound fusion protein travels through the endoplasmic reticulum (ER) and Golgi to the plasma membrane, where the invertase moiety exposed to the periplasm could hydrolyse sucrose, thus allowing growth of the cells. To test this possibility, a ER retrieval signal (DEKKMP) was added to the C-terminus of the invertase-HA-APP (590-695) fusion protein. This signal has been shown to direct retrieval of transmembrane proteins from Golgi to the ER (Gaynor, E.C. et al. (1994) J. Cell. Biol. 127, 653-665). Fig. 6 shows that expression of this novel fusion protein in cells lacking the  $\alpha$ -secretases did not confer them any ability to grow on sucrose plates above that observed upon transformation of an empty plasmid (negative control). However, co-expression of this fusion protein with the  $\alpha$ -secretases restored the ability of the transformed cells to grow on



sucrose plates almost to the extend achieved by expressing wild type invertase.

While there are shown and described presently preferred embodiments of the invention, it is to be distinctly understood that the invention is not limited thereto but may be otherwise variously embodied and practiced within the scope of the following claims.

Claims

1. A method for the identification of one of  
a pair or a group of interacting proteins comprising  
transforming suitable host cells such that  
5 under selective conditions they only survive in the  
presence of desired protein-protein interaction due to  
the release of at least one protein that is essential for  
cell survival under selective conditions, whereby at  
least one of said proteins essential for cell survival is  
10 a secreted protein that in the presence of desired  
interaction is released by cleavage of a fusion protein  
comprising said secreted protein by means of a suitable  
secretase.

2. The method of claim 1 wherein said  
15 proteins are a secretase and a protein comprising a  
cleavage site of said secretase, wherein  
suitable host cells are transformed to  
express under suitable conditions both of said proteins,  
whereby one of said proteins is a candidate  
20 protein of a library to be screened, and whereby  
one of said proteins is present as a fusion  
protein with a secreted protein,  
said host cells transformed with said  
proteins are cultivated under conditions allowing  
25 expression of both proteins, and  
said conditions furthermore allow cell  
survival only in the presence of said secreted protein  
that has been released from said fusion protein due to  
secretase activity exerted on the substrate of said  
30 secretase.

3. The method of claim 1 or 2 wherein said  
interacting protein to be identified is a secretase and  
wherein  
suitable host cells are transformed to  
35 express under suitable conditions a target membrane  
protein as a fusion protein with a secreted protein

said suitable host cells transformed to express said fusion protein are further transformed with a library encoding putative secretases

said host cells transformed with said fusion  
5 protein and said library are cultivated under conditions allowing expression of both said fusion protein and said library

said conditions furthermore allowing cell survival only in the presence of said secreted protein  
10 that has been released from said fusion protein by a secretase encoded by said library.

4. The method of claim 3, wherein the host cell is an eucaryotic cell.

5. The method of claim 3 or 4, wherein the  
15 host cell is a yeast cell.

6. The method of one of claims 3-5, wherein the target protein is a transmembrane protein.

7. The method of one of claims 3-6, wherein the target protein is a type I transmembrane protein.

20 8. The method of one of claims 3-6, wherein the target protein is a type II transmembrane protein.

9. The method of one of claims 3-8, wherein the secreted protein is a protein with invertase activity or functional fragments of a protein with invertase  
25 activity.

10. The method of one of claims 3-9, wherein the secreted protein is a yeast invertase or functional fragments of a yeast invertase.

11. A secretase obtainable by the method of  
30 one of claims 3-10.

12. A protein obtainable by the method of one of claims 3-10 as diagnostic, preventive or therapeutic agent.

13. The method of claim 1 or 2 wherein said  
35 interacting protein to be identified is a substrate of a known secretase and wherein

suitable host cells are transformed to  
express under suitable conditions said secretase

said suitable host cells transformed to  
express said secretase are further transformed with a  
5 library encoding fusion proteins of a secreted protein  
with candidate substrates of said secretase

said host cells transformed with said  
secretase and said library are cultivated under  
conditions allowing expression of both said secretase and  
10 said library

said conditions furthermore allowing cell  
survival only in the presence of said secreted protein  
released from said library encoded fusion protein by said  
secretase.

15 14. The method of claim 13, wherein the host  
cell is an eucaryotic cell.

15. The method of claim 13 or 14, wherein the  
host cell is a yeast cell.

16. The method of one of claims 13-15,  
20 wherein the secreted protein is a protein with invertase  
activity or functional fragments of a protein with  
invertase activity.

17. The method of one of claims 13-16,  
wherein the secreted protein is a yeast invertase or  
25 functional fragments of a yeast invertase.

18. A protein obtainable by the method of one  
of claims 13-17.

19. The protein of claim 18 as diagnostic,  
preventive or therapeutic agent.

30 20. The method of claim 1 wherein said  
interacting protein to be identified is a protein  
interacting with a known target membrane protein and  
wherein

suitable host cells are transformed to  
35 express under suitable conditions said target membrane  
protein as a fusion protein with a secreted protein  
whereby the two moieties of said fusion protein are

linked by a recognition site forming a cleavage site that is cleaved by a defined secretase

said host cells transformed to express said fusion protein are further transformed with a library  
5 encoding fusion proteins of said defined secretase with candidate interaction partner proteins of said target protein

said host cells transformed to express said fusion protein and said library are cultivated under  
10 conditions allowing expression of both said target membrane fusion protein and said library

said conditions furthermore allowing cell survival only in the presence of said secreted protein released from said fusion protein by said defined  
15 secretase due to an interaction of the known target membrane protein with the library encoded interaction partner protein.

21. The method of claim 20, wherein the host cell is a eucaryotic cell.

20 22. The method of claim 20 or 21, wherein the host cell is a yeast cell.

23. The method of one of claims 20-22, wherein the target protein is a transmembrane protein.

24. The method of one of claims 20-23,  
25 wherein the target protein is a type I transmembrane protein.

25. The method of one of claims 20-23, wherein the target protein is a type II transmembrane protein.

30 26. The method of one of claims 20-25, wherein the secreted protein is a protein with invertase activity or functional fragments of a protein with invertase activity.

27. The method of one of claims 20-26,  
35 wherein the secreted protein is a yeast invertase or functional fragments of a yeast invertase.

28. A protein obtainable by the method of one of claims 20-27.

29. The protein of claim 28 as diagnostic, preventive or therapeutic agent.

5 30. The method of claim 1 or 2 wherein said interacting protein to be identified is a secretase and wherein

suitable host cells are transformed to express under suitable conditions a membrane protein as a  
10 fusion protein that comprises said membrane protein, a secreted protein and a transcriptional activator which is part of a reporter system that is stably integrated into the genome of said host cells

said host cells transformed to express said  
15 fusion protein are further transformed with a library encoding candidate secretases

said host cells transformed to express said fusion protein and said library are cultivated under conditions allowing expression of both said fusion  
20 protein and said library

said conditions furthermore allowing cell survival and selection of positive cells only in the presence of a library encoded secretase that cleaves said fusion protein and thus releases the secreted protein and  
25 the transcriptional activator from said fusion protein and thus allows cell survival and selection of positive cells.

31. The method of claim 30, wherein the host cell is a eucaryotic cell.

30 32. The method of claim 30 or 31, wherein the host cell is a yeast cell.

33. The method of one of claims 30-32, wherein said fusion protein comprises a known membrane protein and a transcriptional activator.

35 34. The method of one of claims 30-33, wherein the transcriptional activator is LexA-VP16.

35. The method of one of the claims 30-32, wherein said fusion protein comprises a known membrane protein and a secreted protein.

36. The method of claim 35, wherein said  
5 secreted protein is a protein with invertase activity or functional fragments of a protein with invertase activity.

37. The method of claim 36 wherein said  
secreted protein is a yeast invertase or functional  
10 fragments of a yeast invertase.

38. A secretase obtainable by a method of one of claims 30-37.

39. The secretase of claim 38 as diagnostic, preventive or therapeutic agent.

15 40. The method of claim 1 wherein said interacting protein to be identified is able to process a specific secretase and wherein

suitable host cells are transformed to express under suitable conditions a target protein of  
20 said secretase as a fusion protein with a secreted protein

said host cells transformed to express said fusion protein are further transformed to express an unprocessed specific secretase

25 said host cells transformed to express said fusion protein and said unprocessed secretase are further transformed with a library encoding putative processing proteins of said secretase

said host cells transformed to express said  
30 fusion protein, said unprocessed secretase and said library are cultivated under conditions allowing expression of said fusion protein, said unprocessed secretase and said library

said conditions furthermore allowing cell  
35 survival only in the presence of protein encoded by said library that processes said specific secretase to an

active secretase and thus allows release of said secreted protein from said fusion protein and thus cell survival.

41. The method of claim 40 wherein the host cell is an eucaryotic cell.

5           42. The method of claim 40 or 41 wherein the host cell is a yeast cell.

          43. The method of one of claims 40-42 wherein the secreted protein is a protein with invertase activity or functional fragments of a protein with invertase  
10 activity.

          44. The method of one of claims 40-43 wherein the secreted protein is a yeast invertase or functional fragments of a yeast invertase.

          45. The method of one of claims 40-44 wherein  
15 the membrane protein is a transmembrane protein.

          46. The method of one of claims 40-45 wherein the membrane protein is a type I transmembrane protein.

          47. The method of one of claims 40-45 wherein the membrane protein is a type II transmembrane protein.

20           48. The method of one of claims 40-47 wherein said fusionprotein comprises a target membrane protein, a secreted protein and a transcriptional activator.

          49. A protein obtainable by a method of one of claims 40-49.

25           50. The protein of claim 49 as a diagnostic, preventive or therapeutic agent.



# Fig. 1 Disruption of YAP3 and MKC7

## Yap3p and Mkc7p:

- two related GPI-linked yeast aspartyl proteases
- responsible for  $\alpha$ -secretase-type cleavage of APP in yeast

Zhang et al., BBA 1359 (1997) 110-22, Komano et al., JBC 273 (1998) 31648-51

Fig. 1A






Fig. 1B




# Fig. 1 (continued)

Fig. 1C



Selection on  
-His plates


Fig. 1D



## Fig. 2 Disruption of SUC2


SUC2 knock out in YAP3+ MKC7+ ( $\alpha$ +) and yap3- mkc7- ( $\alpha$ -) genetic background in *S. cerevisiae* strain JPY9

Fig. 2A




Selection on  
-Ura plates

Fig. 2B Genomic PCR controls



# Fig. 3 Genetic test for suc2 disruption

Transformation of a SUC2 expression vector in JPY9  $\alpha$  +/SUC2- and JPY9  $\alpha$  -/SUC2- background



Transformation 24.10.99

# Fig. 4 SUC2-APP fusion with N-terminal truncated APP

Fig. 4A

$\alpha$  - background






Fig. 4B



Transformation 5.11.99


6 / 7

**Fig. 5 SUC2-APP(590-695) is activated in  $\alpha$  + background**



*Transformation 10.11.99*

**Fig. 6 ER retrieval signal in  $\alpha +$  and  $\alpha -$  background**



# INTERNATIONAL SEARCH REPORT

National Application No

PCT/IB 00/00552

**A. CLASSIFICATION OF SUBJECT MATTER**  
 IPC 7 C12N9/64 C07K14/47 C12N9/26

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, WPI Data, PAJ, MEDLINE, CHEM ABS Data, SCISEARCH, BIOTECHNOLOGY ABS, EMBASE

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 13488 A (SCHERING AG ;DYRKS THOMAS (DE); HAERTEL MARION (DE); TURNER JONATH) 2 April 1998 (1998-04-02) examples 1-4	1-10, 13-17, 20-27
Y	BOEHM JOHANNES ET AL: "Kex2-dependent invertase secretion as a tool to study the targeting of transmembrane proteins which are involved in ER fwdarw Golgi transport in yeast." EMBO (EUROPEAN MOLECULAR BIOLOGY ORGANIZATION) JOURNAL, vol. 13, no. 16, 1994, pages 3696-3710, XP000608640 ISSN: 0261-4189 page 3699, left-hand column, last paragraph -right-hand column, paragraph 2; figure 1	1-10, 13-17, 20-27

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*&\* document member of the same patent family

Date of the actual completion of the international search

26 February 2001

Date of mailing of the international search report

23/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

ALCONADA RODRIG., A



# INTERNATIONAL SEARCH REPORT

ational Application No

PCT/IB 00/00552

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DIXON E P ET AL: "AN INVERSE MAMMALIAN TWO-HYBRID SYSTEM FOR BETA SECRETASE AND OTHERPROTEASES" ANALYTICAL BIOCHEMISTRY,US,ACADEMIC PRESS, SAN DIEGO, CA, vol. 249, no. 2, 1997, pages 239-241, XP000914815 ISSN: 0003-2697 the whole document</p>	30-37
A	<p>STEINER H ET AL: "AN IN VIVO ASSAY FOR THE IDENTIFICATION OF TARGET PROTEASES WHICH CLEAVE MEMBRANE-ASSOCIATED SUBSTRATES" FEBS LETTERS,ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM,NL, vol. 463, 17 December 1999 (1999-12-17), pages 245-249, XP000946647 ISSN: 0014-5793 figure 1</p>	30-37
A	<p>DE 198 56 261 C (HOECHST MARION ROUSSEL DE GMBH) 30 March 2000 (2000-03-30) page 3, line 54 -page 5, line 24; examples 1-3</p>	30-37
A	<p>KAISER C A ET AL: "MANY RANDOM SEQUENCES FUNCTIONALLY REPLACE THE SECRETION SIGNAL SEQUENCE OF YEAST INVERTASE" SCIENCE,US,AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE, vol. 235, no. 4786, 16 January 1987 (1987-01-16), pages 312-317, XP000610609 ISSN: 0036-8075 page 312, right-hand column, last paragraph; table 1</p>	1-10, 13-17, 20-27, 30-37, 40-48
A	<p>KOHORN BRUCE D: "Isolation of cDNAs encoding proteases of known specificity using a cleavable GA14 protein." METHODS (ORLANDO), vol. 5, no. 2, 1993, pages 156-160, XP000987043 ISSN: 1046-2023 figures 1,2</p>	1-10, 13-17, 20-27, 30-37, 40-48

-/-

## INTERNATIONAL SEARCH REPORT

national Application No

PCT/IB 00/00552

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>KOMANO HIROTO ET AL: "Involvement of cell surface glycosyl-phosphatidylinositol-linked aspartyl proteases in alpha-secretase-type cleavage and ectodomain solubilization of human Alzheimer beta-amyloid precursor protein in yeast." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 273, no. 48, 27 November 1998 (1998-11-27), pages 31648-31651, XP002161371 ISSN: 0021-9258 the whole document</p>	<p>1-10, 13-17, 20-27, 30-37, 40-48</p>

Continuation of Box I.2

Claims Nos.: 11, 12, 18, 19, 28, 29, 38, 39, 49, 50

Present claims 11, 12, 18, 19, 28, 19, 38, 19, 49 and 50 relate to secretases, secretase-interacting proteins, polypeptides interacting with a known transmembrane protein and a secretase-processing polypeptide defined by reference to a desirable characteristic or property, namely their ability to be detected in the assay as defined in claims 3-10, 13-17, 20-27, 30-37 and 40-49.

The claims cover all polypeptides having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for none of such proteins. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the polypeptides by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has not been carried out for those claims..

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 00/00552

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9813488	A	02-04-1998	DE 19641180 A	26-03-1998
			AU 4775797 A	17-04-1998
DE 19856261	C	30-03-2000	AU 1968000 A	26-06-2000
			WO 0034511 A	15-06-2000

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**